

CCR9 rabbit pAb

Cat No.: ES11459

For research use only

Overview

Product Name CCR9 rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein .

at AA range: 140-220

Specificity CCR9 Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name C-C chemokine receptor type 9 (C-C CKR-9)

(CC-CKR-9) (CCR-9) (G-protein coupled receptor 28)

(GPR-9-6) (CD antigen CDw199)

Gene Name CCR9 GPR28

Cellular localizationCell membrane ; Multi-pass membrane protein .PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 40kD
Human Gene ID 10803
Human Swiss-Prot Number P51686

Alternative Names

Background The protein encoded by this gene is a member of

the beta chemokine receptor family. It is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptors are key regulators of the thymocytes

migration and maturation in normal and

inflammation conditions. The specific ligand of this




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



receptor is CCL25. It has been found that this gene is differentially expressed by T lymphocytes of small intestine and colon, suggested a role in the thymocytes recruitment and development that may permit functional specialization of immune responses in different segment of the gastrointestinal tract. This gene is mapped to the chemokine receptor gene cluster region. Two alternatively spliced transcript variants have been described. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012],

Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950

