

O10D3 rabbit pAb

Cat No.: ES11737

For research use only

Overview

Product Name O10D3 rabbit pAb

Host species Rabbit WB;ELISA **Applications**

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein.

at AA range: 230-310

O10D3 Polyclonal Antibody detects endogenous Specificity

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage** Putative olfactory receptor 10D3 (HTPCRX09) **Protein Name**

(Olfactory receptor OR11-293)

Gene Name OR10D3 OR10D3P

Cell membrane; Multi-pass membrane protein. **Cellular localization** Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml Observed band 34kD

Human Gene ID

Human Swiss-Prot Number

Alternative Names

Q8NH80

Background Olfactory receptors interact with odorant molecules


> in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night



+86-27-59760950