

GLI2 rabbit pAb

Cat No.:ES11962

For research use only

Overview

Product Name	GLI2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from part region of
	human protein
Specificity	GLI2 Polyclonal Antibody detects endogenous levels
	of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathrm{C}$. Avoid repeated freeze-thaw cycles.
Protein Name	Zinc finger protein GLI2 (Tax helper protein)
Gene Name	GLI2 THP
Cellular localization	Nucleus . Cytoplasm . Cell projection, cilium . STK36
	promotes translocation to the nucleus. In
	keratinocytes, it is sequestered in the cytoplasm by
	SUFU. In the absence of SUFU, it translocates to the
	nucleus; [Isoform 1]: Nucleus .; [Isoform 2]:
	Nucleus .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	174kD
Human Gene ID	2736
Human Swiss-Prot Number	P10070
Alternative Names	
Background	This gene encodes a protein which belongs to the
	C2H2-type zinc finger protein subclass of the Gli
	family. Members of this subclass are characterized
	as transcription factors which bind DNA through zinc



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



finger motifs. These motifs contain conserved H-C links. Gli family zinc finger proteins are mediators of Sonic hedgehog (Shh) signaling and they are implicated as potent oncogenes in the embryonal carcinoma cell. The protein encoded by this gene localizes to the cytoplasm and activates patched Drosophila homolog (PTCH) gene expression. It is also thought to play a role during embryogenesis. The encoded protein is associated with several phenotypes- Greig cephalopolysyndactyly syndrome, Pallister-Hall syndrome, preaxial polydactyly type IV, postaxial polydactyly types A1 and B. [provided by RefSeq, Jul 2008],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C