

## VAMP8 rabbit pAb

## Cat No.:ES12390

For research use only

## Overview

| Product Name             | VAMP8 rabbit pAb   |
|--------------------------|--|
| Host species             | Rabbit   |
| Applications             | IHC;IF   |
| Species Cross-Reactivity | Human; Mouse;Rat   |
| Recommended dilutions    | IHC-p 1: 50-200  |
| Immunogen                | Synthesized peptide derived from human VAMP8 AA range: 46-96   |
| Specificity              | This antibody detects endogenous levels of VAMP8<br>at Human/Mouse/Rat   |
| Formulation              | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Storage                  | Store at -20°C. Avoid repeated freeze-thaw cycles.   |
| Protein Name             | VAMP-8   |
| Gene Name                | VAMP8  |
| Cellular localization    | Lysosome membrane ; Single-pass type IV<br>membrane protein . Early endosome membrane ;<br>Single-pass type IV membrane protein . Late<br>endosome membrane ; Single-pass type IV<br>membrane protein . Cell membrane ; Single-pass<br>type IV membrane protein . Zymogen gran |
| Purification             | The antibody was affinity-purified from rabbit<br>antiserum by affinity-chromatography using<br>epitope-specific immunogen.  |
| Clonality                | Polyclonal   |
| Concentration            | 1 mg/ml  |
| Observed band            |  |
| Human Gene ID            | 8673   |
| Human Swiss-Prot Number  | Q9BV40   |
| Alternative Names        | llin.  |
| Background               | This gene encodes an integral membrane protein<br>that belongs to the<br>synaptobrevin/vesicle-associated membrane protein<br>subfamily of soluble N-ethylmaleimide-sensitive  |



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



factor attachment protein receptors (SNAREs). The encoded protein is involved in the fusion of synaptic vesicles with the presynaptic membrane.[provided by RefSeq, Jun 2010],

Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C