

UD11 rabbit pAb

Cat No.: ES12436

For research use only

Overview

Product Name UD11 rabbit pAb

Host species Rabbit
Applications WB

Species Cross-Reactivity Human; Mouse; Rat Recommended dilutions WB 1: 500-2000

Immunogen Synthesized peptide derived from human UD11 AA

range: 36-86

Specificity This antibody detects endogenous levels of UD11 at

Human/Mouse/Rat

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name UD11

Gene Name UGT1A1 GNT1 UGT1

Cellular localization Endoplasmic reticulum membrane ; Single-pass

membrane protein. Cytoplasm, perinuclear region.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

Observed band

Human Gene ID 54658 Human Swiss-Prot Number P22309

Alternative Names

Background This gene encodes a UDP-glucuronosyltransferase,

an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by

+86-27-59760950 ELKbio@ELKbiotech.com


www.elkbiotech.com



are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. The preferred substrate of this enzyme is bilirubin, although it also has moderate activity with simple phenols, flavones, and C18 steroids. Mutations in this gene result in Crigler-Najjar syndromes types I and II and in Gilbert syndrome. [provided by RefSeq, Jul 2008],

four common exons. Four of the alternate first exons

Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950