

Histone H3 (Acetyl Lys56) rabbit pAb

Cat No.:ES20520

For research use only

Overview

Produc t Name	Histone H3 (Acetyl Lys56) rabbit pAb
Host species	Rabbit
Applica tions	WB
Species Cross-R eactivit	Human;Mouse;Rat
y Recom mende d dilution	WB: 1:500-1000
S	Current atia Dautida of Uiston a UD (Acatul Luc C)
lmmun ogen	Synthetic Peptide of Histone H3 (Acetyl Lys56)
Specific ity	The antibody detects endogenous Histone H3 (Acetyl Lys56) protein.
Formul ation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Histone H3.1/Histone H3.2/Histone H3.3
Gene Name	HIST1H3A/HIST1H3B/HIST1H3C/HIST1H3D/HIST1H3E/HIST1H3F/HIST1H3G /HIST1H3H/HIST1H3I/HIST1H3J/HIST2H3A/HIST2H3C/HIST2H3D/H3F3A/H 3F3B
Cellular localiza tion	Nucleus. Chromosome.
Purifica tion Clonalit	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Y	
Concen	
tration	
Observ	15-17kD
ed	
band	
Human	8350/8351/8352/8353/8354/8355/8356/8357/8358/8968
Gene	
ID	
Human	P68431/Q71DI3/P84243
Swiss-P	
rot	
Numbe	
r	
Alterna	HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB;
tive	HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK;
Names	HIST1H3I; H3FF; HIST1H3J; H3FJ; Histone H3.1; Histone H3/a; Histone
	H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/
Backgr	Histones are basic nuclear proteins that are responsible for the
ound	nucleosome structure of the chromosomal fiber in eukaryotes. This
	structure consists of approximately 146 bp of DNA wrapped around a
	nucleosome, an octamer composed of pairs of each of the four core
	histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted
	through the interaction of a linker histone, H1, with the DNA between the
	nucleosomes to form higher order chromatin structures. This gene is
	intronless and encodes a replication-dependent histone that is a member
	of the histone H3 family. Transcripts from this gene lack polyA tails;
	instead, they contain a palindromic termination element. This gene is
	found in the large histone gene cluster on chromosome 6p22-p21.3.
	[provided by RefSeq, Aug 2015],



Western blot analysis of extracts from Hela cells, untreated (-) or treated, 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C