

SESN3 rabbit pAb

Cat No.:ES13146

For research use only

Overview

| Product Name | SESN3 rabbit pAb |
|------------------------------|--|
| Host species | Rabbit |
| Applications | WB;ELISA;IHC |
| Species Cross-Reactivity | Human; Mouse |
| Recommended dilutions | WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 |
| Immunogen | Synthesized peptide derived from human SESN3 AA |
| | range: 35-85 |
| Specificity | This antibody detects endogenous levels of SESN3 at |
| | Human/Mouse |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | SESN3 |
| Gene Name | SESN3 SEST3 |
| Cellular localization | Cytoplasm . |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | |
| Human Gene ID | 143686 |
| Human Swiss-Prot Number | P58005 |
| Alternative Names | |
| Background | This gene encodes a member of the sestrin family of |
| | stress-induced proteins. The encoded protein |
| | reduces the levels of intracellular reactive oxygen |
| | species induced by activated Ras downstream of |
| | RAC-alpha serine/threonine-protein kinase (Akt) and |
| | FoxO transcription factor. The protein is required for |
| | normal regulation of blood glucose, insulin |
| | resistance and plays a role in lipid storage in obesity. |
| | Alternative splicing results in multiple transcript |
| | |




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





variants. [provided by RefSeq, Oct 2012],

Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C